

SREE CHITRA TIRUNAL INSTITUTE FOR MEDICAL SCIENCES & TECHNOLOGY THIRUVANANTHAPURAM—695 011, INDIA.

(An Institute of National Importance under Govt.of India)
Phone—(91)0471—2443152 Fax—(91)0471—2446433, 2550728
Email-sct@sctimst.ac.in Web site—www.sctimst.ac.in

ROLL NUMBER	

WRITTEN TEST FOR THE POST OF TECH.ASST. ANAESTHESIA - A To B

DATE: 19/03/2024

TIME: 9.30 to 10.30 AM

DURATION: 60 MINUTES

Total Marks: 50

INSTRUCTIONS TO THE CANDIDATES

- 1. Write your Roll Number on the top of the Question Booklet and in the answer sheet.
- 2. Each question carries 1 mark.
- 3. There will not be any Negative Marking.
- 4. Write legibly the alphabet of the most appropriate answer (A, B, C or D) in the separate answer sheet provided.
- 5. Over-writing is not permitted.
- 6. Candidate should sign in the question paper and answer sheet.
- 7. No clarifications will be given.
- 8. Candidate should hand over the answer sheet to the invigilator before leaving the examination hall.

Signature of the Candidate

land 31mg

MFCP-1 TECHNICIAL ASSISTANT ANAESTHESIA

1. Whi	ch of the following colouring agents is not used	for indi	
a.	Mimosa Z	c.	Ethyl violet
b.	Phenolphthalein	d.	Methylene blue
2. Whi	ch of the following metals is compatible with th	e Magn	etic resonance Imaging (MRI)?
a.	Ferritic steel	c.	Nickel alloy
b.	Gold	d.	Cobalt alloy
	ch one of the following is a complication of sitt	ing posit	tion in patients undergoing posterior
a.	Venous air embolism	C.	Tension pneumocephalus
b.	Quadriparesis	d.	All of the above
4. Gado	olinium is avoided in patients with		
· a.	History of bronchial asthma	c.	History of renal failure
b.	History of heart failure	d.	History of liver failure
5 Whie	ch of the following inhalational agents is pleasa	nt during	a anaesthesia industion in Children?
a.	Sevoflurane		Desflurane
L .	Isoflurane		Enflurane
D.	isoriurane	u.	Emurane
6. Which	ch statement regarding blood component transfu	ision is v	wrong?
a.	Recipient of AB group cannot be given plasma	of B gr	oup donor
b.	O negative group is considered universal dono	r	
c.	Donor of B negative group cannot donate bloo		negative group
d.	Recipient of O group cannot receive plasma of		
7. The	cause of reddish discolouration of urine following	ng mass	ive blood transfusion may be because
a.	Hemophilia	c.	Hemoglobinuria
b.	hyperbilirubinemia		Rhabdomyolysis
0.	nyperonina omonia.	u.	Riaddoniyorysis
2.0	atrial line placed during weaning a patient from	cardiop	ulmonary bypass operated for arterial
	surgery is for		
a.	Measuring cardiac filling pressures		Assessing systemic vascular resistance
b.	Estimating pulmonary hypertension	d.	Measuring cardiac output
9. The	procedure recommended by the Difficult airway	algorith	nm in a case of anticipated difficult
	entilation and intubation in adults is	0	P
a.	Awake fibreoptic intubation	C	Awake direct laryngoscopy
	Awake video laryngoscopic intubation		All of the above
, J.	Tivake video kilyngoscopie intaoation	u.	All of the above
10. The	Laryngeal mask airway of choice in a 2-year-o	ld child	weighing 12 kg is
a.	1	c.	2
b.	1.5	d.	2.5
11. The	value of BIS denoting adequate depth of anaest	thesia du	aring cardiac surgery is
a.	40-50		80-90
ĥ	10-20	d	00 100

	a. b. c.	Patients with upper lower limb muscle weakness due to motor neuropathy Patients with chronic kidney disease Patients with severe abdominal infections				
			ardi	ac failure		
13.	a.b.c.	Pressure regulated volume control ventilation Airway pressure release ventilation Synchronous intermittent mandatory ventilation- Pressure support ventilation				
14.	Sug	gammadex is ideally used to reverse the effect of				
	a.	Succinyl choline	C.	Midazolam		
	b.	Rocuronium	d.	Ketamine		
15.		e Target Controlled Infusion (TCI) models of propo				
	a.	Marsh		Kataria		
	b.	Schnider	d.	Minto		
	he c	nich of the regional blocks administered during adul	t ca	rdiac surgery is not in the anterior part		
		Transverse thoracis plane block		Pectoralis nerve block (PECS 2)		
	b.	Pecto-intercostal fascial plane block	d.	Erector spinae plane block		
17.		ile administering total intravenous anaesthesia which				
		Entropy monitor		Neuromuscular monitoring		
	D.	NIRS	d.	Evoked potential monitoring		
		ich of the following errors can be produced by pres gas sample?	enc	e of excess of heparin in the arterial		
	a.	Low PaO2	c.	Metabolic alkalosis		
	b.	High PaCO2	d.	Metabolic acidosis		
19.	Car	diopulmonary resuscitation algorithm for adults fol	low	ing asystole includes		
	a.	Early defibrillation	c.	Administration of atropine		
	b.	Administration of adrenaline	d.	All of the above		
20.	Mat	terial of the Microcuff of Microcuff endotracheal tu				
		Microthin silicone		Microthin polyurathane		
	b.	Microthin rubber	d.	None of the above		
21.		Partial pressure of Oxygen (PaO2) corresponding to 90 mm Hg		n SpO2 value of 90% is 60 mm Hg		
		85mm Hg		50 mm Hg		
	_	oshini rig	u.	30 mm rig		
22.	Sele	ect incorrect statement on low flow anaesthesia				
		Carbon dioxide should be removed during low flow	v an	esthesia		
		FiO2 monitoring is absolutely necessary				
		Fresh gas flow should exceed alveolar ventilation				
		Usually 1 to 2 litres of fresh gas flow is delivered				

12. Succinyl choline can be safely used in which of the following conditions?

- 23. American society of Anesthesiology (ASA) class 2 means
 - a. A moribund patient who is not expected to survive without operation
 - b. A patient with mild systemic disease
 - c. A declared brain-dead patient
 - d. A patient with recent myocardial infarction or cerebrovascular accident, which is a constant threat to the life
- 24. Select a correct statement: A preoperative ABG estimation can
 - a. Act as baseline/benchmark value for perioperative blood gases
 - b. Can quantify the impairment of gas exchange
 - c. Can detect CO2 retention
 - d. All of the above
- 25. An appropriate preoperative strategy for a neonate scheduled for arterial switch operation would be
 - a. Breast feed 4 hours before surgery
 - b. Breast feed 6 hours before surgery
 - c. Intramuscular morphine 1 hour before surgery
 - d. Nasal midazolam drops 2 hours before surgery
- 26. All are true about benzodiazepines except
 - a. They have convulsant properties
 - b. They have anesthetic properties

- c. They give freedom from anxiety
- d. They can be given orally
- 27. Select correct statement about Glycopyrolate:
 - a. Does not cause central anticholinergic effects
 - b. Results in less tachycardia than atropine
 - c. More longer acting than atropine in drying salivary secretions
 - d. All of the above
- 28. Select wrong statement about nitric oxide
 - a. It is administered via inhalational route
 - b. It is used in the treatment of pulmonary hypertension
 - c. Severe hypotension due reduction in systemic vascular resistance is its adverse effect
 - d. Average therapeutic concentration used is about 20 ppm.
- 29. Identify wrong statement about medical gas cylinders
 - a. Oxygen is stored in liquid form in type D cylinders mounted on anesthesia workstations
 - b. Carbon dioxide cylinders are Grey in colour
 - c. Pin index for mounting of nitrous oxide cylinder on anesthesia workstation is 3-5
 - d. Type H cylinder is bigger than type E
- 30. Capacity of oxygen cylinder size E is
 - a. 680L

c. 100L

b. 50L

d. 150L

- 31. Select a wrong statement about intra-aortic balloon pump
 - a. Balloon catheter has a channel to monitor invasive aortic pressure
 - b. IABP inflation and deflation can be triggered using ECG
 - c. IABP balloon can be inserted via axillary artery
 - d. Balloon is inflated using carbon dioxide gas
- 32. Select a CORRECT statement about cardiac output monitoring
 - Thermodilution cardiac output is calculated by increase in temperature measured at pulmonary artery
 - b. PICCO is a non-invasive type of cardiac output measurement method
 - c. Pulse contour analysis method will be reliable in the presence of IABP counter pulsation
 - d. Cardiac output can be measured using transesophageal echocardiography
- 33. Select appropriate answer: Anesthetic gas scavenging system
 - a. Recirculates exhaled gas

b. Absorbs CO2

- Transports waste anaesthetic gases to the outside
- d. None of the above
- 34. Identify WRONG statement regarding transesophageal echocardiography
 - a. Thermal heating is much more with a 3DTEE probe than a 2DTEE probe.
 - b. Electrocautery can degrade the echocardiographic image
 - c. TEE probe of one system (iE33, Philips Ultrasound) cannot be fitted on to other system (GE Vivid E 95 ultrasound)
 - d. 3DTEE probe can be used for echocardiographic examination in infants
- 35. The modern autoclave systems

00

- a. Uses steam at 134 degree Celsius at 2 bar pressure for 3.5 minutes
- b. Steam is evacuated and replaced by sterile air
- c. The entire cycle takes about 10 minutes
- d. All of the above
- 36. Select a wrong statement about fiberoptic bronchoscopy during lung isolation
 - a. It is held in left hand and tip control knob is operated using thumb by a right-handed person
 - b. Right bronchus is shorter than the left bronchus
 - c. Bronchial cuff of a correctly placed DLT should not be seen near carina when FOB is advanced through tracheal lumen
 - d. Two cuffs of EZ blocker should be seen placed in right and left bronchus
- 37. Which among the following modalities can be useful in monitoring both the respiratory and circulatory systems during conduct of CPR?

a. Non-invasive blood pressure

c. Capnography

b. ECG

d. Airway pressure

- 38. Which of the following agent causes 'dissociative anesthesia'?
 - a. Thiopentone

c. Etomidate

b. Ketamine

d. Propofol

- 39. Which is the drug of choice in managing anaphylaxis?
 - a. Epinephrine

c. Norepinephrine

b. Dobutamine

d. Vasopressin

40. The vocal cords are supplied by which nerv	ve?
a. Glossopharyngeal	c. Hypoglossal
b. Vagus	d. Facial
41. During airway examination, the Anesthesic Mallampatti class the patient belongs?	ologist is not able to see the soft palate. Which
a. Class 1	c. Class 3
b. Class 2	d. Class 4
42. Which iv fluid is not ideal in neuroanaesthe	esia
 a. Glucose containing iv fluids 	c. Ringer lactate
b. Plasmalyte	d. Normal saline
43. Which anesthetic agent would be preferred	in severe asthma patient?
a. Thiopentone	c. Etomidate
b. Ketamine	d. Propofol
44. Which volatile agent can cause seizures at l	nigh doses?
a. Sevoflurane	c. Desflurane
b. Isoflurane	d. Enflurane
45. In lung protective ventilation, the tidal volu	ime set on ventilator should be
a. 3-5	c. 10-12
b. 6-8	d. 12-14
46. The problem with the use of NSAIDs like of	liclofenac in neurosurgeries is
a. Poor analgesia	c. Irritant to veins
b. Delayed recovery	d. Potential to increase bleeding
47. Which of following nerve blocks is NOT pe	erformed during awake intubation?
 a. Glossopharyngeal nerve block 	 c. Retro laryngeal nerve block
b. Superior laryngeal nerve block	d. Trans-laryngeal block
48. Target control infusion pumps are used for	administering
a. Propofol	c. Atracurium
b. Fentanyl	d. Noradrenaline
	on of aortic cross-clamp during thoracoabdomina
aortic aneurysm, following is done except:	
 a. Spinal CSF drainage to keep CSF press 	
b. Administration of sodium pentothal, ste	
c. Re-implantation of retrograde bleeding	intercostal arteries below cross-clamp
d. Administration of sodium nitroprusside	to control proximal hypertension
	vere cyanotic spell during dissection around heart
before establishing CPB. Following can be adm	ninistered, except
a. Sodium bicarbonate	c. Isoproterenol infusion
b. 100% oxygen inhalation	d. Phenylephrine bolus

Technician Anaesthesia

Answer Keys

	1. D			27. D
	2. B			28. C
	3. D			29. A
	4. C			30. A
	5. A			31. D
	6. D			32. D
	7. C			33. C
	8. A			34. D
	9. D			35. D
	10. C			36. C
	11. A			37. C
	12. D			38. B
	13. B			39. A
	14. B			40. B
	15. D			41. D
	16. D	540		42. A
	17. A			43. B
	18. D			44. D
	19. B		e	45. B
6	20. C			46. D
	21. C			47. C
	22. C			48. A
	23. B			49. D
	24. D			50: C

25. A

26. A

Kant 19/3/24